

HIGHER GENUS MEAN CURVATURE 1 CATENOIDS IN HYPERBOLIC AND DE SITTER 3-SPACES

SHOICHI FUJIMORI AND WAYNE ROSSMAN

ABSTRACT. We show existence of constant mean curvature 1 surfaces in both hyperbolic 3-space and de Sitter 3-space with two complete embedded ends and any positive genus up to genus twenty. We also find another such family of surfaces in de Sitter 3-space, but with a different non-embedded end behavior.

INTRODUCTION

This paper extends the result in [RS] by K. Sato and the second author, and also the result in [F2] by the first author.

In [RS], it was shown that, although the only complete connected finite-total-curvature minimal immersions in $\mathbf{R}^3$ with two embedded ends are catenoids (Schoen [S]), there do exist complete connected immersed constant mean curvature (CMC) 1 surfaces with two ends in hyperbolic 3-space H^3 that are not surfaces of revolution, although such non-rotational surfaces in H^3 cannot be embedded (Levitt and Rosenberg [LR]). The examples found in [RS] are of genus one, but there exist examples of genus zero as well, called warped catenoid cousins, which we comment on later in this introduction. This comparison is of interest, because minimal surfaces in $\mathbf{R}^3$ and CMC 1 surfaces in H^3 are Lawson correspondents, and therefore have a very close relationship [B], [UY1], [UY2], [RUY1].

In [F2], analogous spacelike surfaces in de Sitter 3-space S_1^3 were shown to exist. Likewise, in this non-Riemannian situation, there is a similar close relationship between spacelike maximal surfaces in Minkowski 3-space $\mathbf{R}_1^3$ and spacelike CMC 1 surfaces in S_1^3 . The interest in these surfaces stems largely from the nature of their singular sets.

There is a well-known classical Weierstrass representation for minimal surfaces in $\mathbf{R}^3$, and a very similar Weierstrass type representation for maximal surfaces in $\mathbf{R}_1^3$ ([K], [UY4] for example). Because of the relationships described above, we have again Weierstrass type representations for CMC 1 surfaces in H^3 ([B], [UY1] for example) and for CMC 1 surfaces in S_1^3 ([AA], [F1], [FRUY]). These representations are used here, for H^3 and S_1^3 , in Equations (1.2) and (1.3). Furthermore, because of all of these relationships, the Osserman inequality for minimal surfaces in $\mathbf{R}^3$ has analogs for maximal surfaces in $\mathbf{R}_1^3$ ([UY4]), and CMC 1 surfaces in H^3 ([UY3]) and S_1^3 ([F1], [FRUY]).

The examples found in [RS] and [F2] were only of genus 1, and the purpose in this article is to show:

- (1) the method of [RS] can be extended to give examples of any positive genus up to genus twenty, and probably any even higher genus as well, without requiring a multi-dimensional period problem (showing that a simplification of a comment made in the introduction of [RS] is possible), and

Date: February 25, 2010.

2000 Mathematics Subject Classification. Primary 53A10; Secondary 65D17.

Key words and phrases. constant mean curvature 1 surface, higher genus surface, hyperbolic 3-space, de Sitter 3-space.

- (2) in light of recent work on CMC 1 surfaces with singularities in S_1^3 , the same method will give CMC 1 surfaces of any genus (at least up to genus twenty) and two embedded ends in S_1^3 , and
- (3) although the CMC 1 surfaces in H^3 and S_1^3 have a similar mathematical construction, the behavior of the ends of the surfaces in S_1^3 is more complicated to analyze, related to the fact that the group $SU(1, 1)$ used in the S_1^3 case is not compact (although $SU(2)$, used in H^3 , is). To demonstrate this, we find a family of surfaces in S_1^3 with hyperbolic ends (the term “hyperbolic ends” was defined in [F1] and [FRUY]).

CMC 1 surfaces with certain kinds of singularities in S_1^3 were called CMC 1 faces in [F1] and [FRUY]. Regarding the third point above, in [F1] and [FRUY] it was shown that ends of CMC 1 faces in S_1^3 come in three types: elliptic, hyperbolic and parabolic. However, ends of CMC 1 surfaces in H^3 will always be elliptic. Because of this, in the S_1^3 case an extra argument is needed to demonstrate the numerical result just below, and we give that argument at the end of this paper. Our main result is this:

Numerical result: There exists a one-parameter family of CMC 1 genus k complete properly immersed surfaces in H^3 with two embedded ends, for any positive integer $k \leq 20$. Likewise, again for any positive integer $k \leq 20$, there exist two one-parameter families of genus k CMC 1 faces in S_1^3 , one with two complete embedded elliptic ends, and one with two weakly complete (in the sense of [FRUY]) hyperbolic ends.

We expect the result is true for many integers $k \geq 21$ as well, if not all integers $k \geq 21$.

Note that the surfaces in H^3 and the first family of surfaces in S_1^3 have the nice property that they are embedded outside of a compact set.

Here we are interested in the case that k is positive, but there do exist CMC 1 surfaces with genus 0 and embedded ends that are not surfaces of revolution. In the H^3 case, they can be found in [UY1] (Theorem 6.2) and [RUY2] (where they are called warped catenoid cousins), and those surfaces in H^3 imply the existence of corresponding non-rotational examples in S_1^3 by Theorem 5.6 in [F1].

The surfaces in the above numerical result are not known to exist by any rigorous mathematical method, so, like in [RS] and [F2], we rely on numerics at one step to show this result. In particular, we show numerically that a certain continuous function from the real line to the real line is positive at one point and negative at another, thus implying by the intermediate value theorem that it has a zero.

We provide some graphics of higher genus catenoids in H^3 , see Figure 1. (Because the surfaces in S_1^3 have singularities, making them more difficult to visualize globally, the computer graphics become less helpful in this case, and we do not show such graphics here.)

We conclude this introduction with two related remarks:

- (1) Although there do not exist any genus 1 complete connected finite-total-curvature minimal immersions in $\mathbf{R}^3$ with two embedded ends, there do exist genus 1 maximal surfaces (they can actually be complete maxfaces in the sense of [UY4]) with two embedded ends in $\mathbf{R}_1^3$. See Kim-Yang [KY].
- (2) If one allows the ends to be non-embedded, there do exist examples of complete connected finite-total-curvature minimal surfaces with two ends and positive genus. See Fujimori-Shoda [FS] for example.

Acknowledgements. The authors are very grateful to Masaaki Umehara, Kotaro Yamada and Seong-Deog Yang for many fruitful discussions, without which the authors would not have found the result here.




FIGURE 1. Half cut-away of higher genus catenoids in H^3 (in the Poincaré ball model), on the left, and central portions of those same surfaces on the right, for $k = 1$, $k = 2$, $k = 3$ and $k = 8$.

1. THE WEIERSTRASS DATA

Here we use a more general Weierstrass data than in [RS] and [F2], allowing the genus to be any positive number.

Take the compact Riemann surface

$$\overline{M} = \left\{ (z, w) \in (\mathbf{C} \cup \{\infty\})^2 \mid w^{k+1} = z \left(\frac{z - \lambda^{-1}}{\lambda - z} \right)^k \right\},$$

where k is any positive integer and λ is any real constant such that $\lambda > 1$. This $\overline{M}$ has the structure of a Riemann surface, and z provides a local complex coordinate for $\overline{M}$ at all but four points. At those four points $(0, 0)$, (∞, ∞) , (λ, ∞) and $(\lambda^{-1}, 0)$, we can take a local coordinate ζ_0 , ζ_∞ , ζ_λ and $\zeta_{\lambda^{-1}}$ satisfying $\zeta_0^{k+1} = z$, $\zeta_\infty^{-k-1} = z$, $\zeta_\lambda^{k+1} = z - \lambda$ and $\zeta_{\lambda^{-1}}^{k+1} = z - \lambda^{-1}$, respectively. By applying the Riemann-Hurwitz relation, we find that this Riemann surface has genus k . Then take

$$M = \overline{M} \setminus \{(0, 0), (\infty, \infty)\}.$$

Here $(0, 0)$ and (∞, ∞) will represent the two ends of the surfaces we will construct (surfaces having domain M). Let $\widetilde{M}$ be the universal cover of M .

We now take the Weierstrass data

$$(1.1) \quad G = \lambda^{k/(k+1)} w, \quad \Omega = c \cdot \frac{dz}{zw}.$$

Here c is any nonzero real constant.

Remark 1.1. Multiplying the hyperbolic Gauss map G by a constant is equivalent to just a rigid motion of the surface, in both H^3 and S_1^3 . So we could have chosen G to be $G = w$ in Equation (1.1), as our goal is to construct surfaces in H^3 and S_1^3 . However, when considering relations with minimal surfaces in $\mathbf{R}^3$ and maximal surface in $\mathbf{R}_1^3$, the choice of G in (1.1) will prove to be useful, as we will see in Remark 2.4. So here we use the hyperbolic Gauss map G as given in (1.1).

Note that $\deg(G) = k + 1$. Now take a solution

$$F : \widetilde{M} \rightarrow \mathrm{SL}(2, \mathbf{C})$$

of

$$(1.2) \quad dF \cdot F^{-1} = \begin{pmatrix} G & -G^2 \\ 1 & -G \end{pmatrix} \Omega.$$

Noting that, with $a^* = \bar{a}^t$,

$$H^3 = \{aa^* \mid a \in \mathrm{SL}(2, \mathbf{C})\}, \quad S_1^3 = \{ae_3a^* \mid a \in \mathrm{SL}(2, \mathbf{C})\}, \quad e_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix},$$

define

$$(1.3) \quad f_H := FF^* : \widetilde{M} \rightarrow H^3 \quad \text{and} \quad f_S := Fe_3F^* : \widetilde{M} \rightarrow S_1^3.$$

Then f_H (resp. f_S) gives a CMC 1 immersion in H^3 (resp. a CMC 1 face in S_1^3), since

$$(1.4) \quad (1 + |G|^2)^2 \Omega \overline{\Omega}$$

gives a positive definite metric on M , see [UY3] and Theorem 1.9 of [F1]. Note that the Hopf differential of both f_H and f_S is written as

$$Q = \Omega dG = \frac{c\lambda^{k/(k+1)}}{k+1} \cdot \frac{z^2 + \{(k-1)\lambda^{-1} - (k+1)\lambda\}z + 1}{z^2(z - \lambda^{-1})(z - \lambda)} dz^2.$$

2. SYMMETRIES OF THE SURFACE

We define

$$w_0 := \lambda^{-k/(k+1)} \in \mathbf{R} \quad \text{and} \quad \Lambda := e^{2\pi i/(k+1)}.$$

Then it is easy to see that

$$(1, \Lambda^j w_0) \in M \quad \text{for any } j = 0, 1, \dots, k.$$

Consider the symmetries

$$\kappa_1(z, w) = (\bar{z}, \bar{w}), \quad \kappa_2(z, w) = \left(\frac{1}{z}, \frac{1}{\lambda^{2k/(k+1)} w} \right), \quad \kappa_3(z, w) = (\bar{z}, \Lambda \bar{w})$$

on $\overline{M}$. Then we have the following, which follows from a proof analogous to proofs found in [RS] and [F2]:

Lemma 2.1. *Let*

$$F(z, w) = \begin{pmatrix} A & B \\ C & D \end{pmatrix}$$

be a solution of (1.2) with the initial condition $F(1, w_0) = \text{id}$. Then

$$(2.1) \quad F(\kappa_1(z, w)) = \begin{pmatrix} \bar{A} & \bar{B} \\ \bar{C} & \bar{D} \end{pmatrix}, \quad F(\kappa_2(z, w)) = \begin{pmatrix} D & C \\ B & A \end{pmatrix}, \quad F(\kappa_3(z, w)) = \begin{pmatrix} \bar{A} & \Lambda \bar{B} \\ \Lambda^{-1} \bar{C} & \bar{D} \end{pmatrix}.$$

Proof. Note that we have the following relations under the symmetries κ_j :

$$G \circ \kappa_1 = \bar{G}, \quad G \circ \kappa_2 = G^{-1}, \quad G \circ \kappa_3 = \Lambda \bar{G}, \\ \kappa_1^* \Omega = \bar{\Omega}, \quad \kappa_2^* \Omega = -\lambda^{2k/(k+1)} w^2 \Omega, \quad \kappa_3^* \Omega = \Lambda^{-1} \bar{\Omega}.$$

It follows that

$$\begin{aligned} \kappa_1^* \begin{pmatrix} G & -G^2 \\ 1 & -G \end{pmatrix} \Omega &= \overline{\begin{pmatrix} G & -G^2 \\ 1 & -G \end{pmatrix} \Omega}, \\ \kappa_2^* \begin{pmatrix} G & -G^2 \\ 1 & -G \end{pmatrix} \Omega &= \begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix}^{-1} \begin{pmatrix} G & -G^2 \\ 1 & -G \end{pmatrix} \Omega \begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix}, \\ \kappa_3^* \begin{pmatrix} G & -G^2 \\ 1 & -G \end{pmatrix} \Omega &= \begin{pmatrix} \sqrt{\Lambda} & 0 \\ 0 & 1/\sqrt{\Lambda} \end{pmatrix} \overline{\begin{pmatrix} G & -G^2 \\ 1 & -G \end{pmatrix} \Omega} \begin{pmatrix} 1/\sqrt{\Lambda} & 0 \\ 0 & \sqrt{\Lambda} \end{pmatrix}. \end{aligned}$$

Because the initial condition $F(1, w_0) = \text{id}$ satisfies

$$\begin{aligned} \overline{F(1, w_0)} &= F(1, w_0), \quad \begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix}^{-1} F(1, w_0) \begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix} = F(1, w_0), \\ \begin{pmatrix} \sqrt{\Lambda} & 0 \\ 0 & 1/\sqrt{\Lambda} \end{pmatrix} \overline{F(1, w_0)} \begin{pmatrix} 1/\sqrt{\Lambda} & 0 \\ 0 & \sqrt{\Lambda} \end{pmatrix} &= F(1, w_0), \end{aligned}$$

the lemma follows. $\square$

We now consider two loops in M (see Figure 2):

- The loop $\gamma_1 : [0, 1] \rightarrow M$ starts at $\gamma_1(0) = (1, w_0) \in M$. Its first portion has z coordinate in $\{\text{Im}(z) < 0\}$ and ends at a point (z, w) where $z \in \mathbf{R}$ and $0 < z < \lambda^{-1}$. Its second portion starts at (z, w) and ends at $(1, \Lambda w_0)$ and has z coordinate in $\{\text{Im}(z) > 0\}$. Its third portion starts at $(1, \Lambda w_0)$ and ends at $(1/z, \Lambda/(\lambda^{2k/(k+1)} w))$ and has z coordinate in $\{\text{Im}(z) > 0\}$. Its fourth and last portion starts at $(1/z, \Lambda/(\lambda^{2k/(k+1)} w))$ and returns to the base point $\gamma_1(1) = (1, w_0)$ and has z coordinate in $\{\text{Im}(z) < 0\}$.
- The loop $\gamma_2 : [0, 1] \rightarrow M$ starts at $\gamma_2(0) = (1, w_0)$. Its first portion has z coordinate in $\{\text{Im}(z) < 0\}$ and ends at a point (z, w) where $z \in \mathbf{R}$ and $z < 0$. Its second and last portion starts at (z, w) and returns to $\gamma_2(1) = (1, w_0)$ and has z coordinate in $\{\text{Im}(z) > 0\}$.




FIGURE 2. Projection to the z -plane of the loops γ_1 and γ_2 . (Note that $z = 1$ is not a branch point of M , and γ_1 starts at one lift of $z = 1$, and then passes through another lift of $z = 1$, and then returns to the first lift of $z = 1$.)

We will also consider the following two paths (not loops) in M (see Figure 3):




FIGURE 3. Projection to the z -plane of the curves c_1 and c_2 .

- Let $c_1 : [0, 1] \rightarrow M$ be a curve starting at $c_1(0) = (1, w_0)$ whose projection to the z -plane is an embedded curve in $\{\text{Im}(z) < 0\}$, and whose endpoint $c_1(1)$ has a z coordinate so that $z \in \mathbf{R}$ and $0 < z < \lambda^{-1}$.
- Let $c_2(t) : [0, 1] \rightarrow M$ be a curve starting at $c_2(0) = (1, w_0)$ whose projection to the z -plane is an embedded curve in $\{\text{Im}(z) < 0\}$, and whose endpoint $c_2(1)$ has a z coordinate so that $z \in \mathbf{R}$ and $z < 0$.

With $F(1, w_0) = \text{id}$, we solve Equation (1.2) along these two paths to find

$$F(c_1(1)) = \begin{pmatrix} A_1 & B_1 \\ C_1 & D_1 \end{pmatrix}, \quad \text{and} \quad F(c_2(1)) = \begin{pmatrix} A_2 & B_2 \\ C_2 & D_2 \end{pmatrix}.$$

Let τ_j be the deck transformation of $\widetilde{M}$ associated to the homotopy class of γ_j ($j = 1, 2$).

Lemma 2.2. *We have that $F \circ \tau_1 = F\Phi_1$ and $F \circ \tau_2 = F\Phi_2$, where*

$$\begin{aligned} \Phi_1 &:= \begin{pmatrix} \bar{A}_1 & -\bar{C}_1 \\ -\bar{B}_1 & \bar{D}_1 \end{pmatrix} \begin{pmatrix} D_1 & \Lambda C_1 \\ \Lambda^{-1} B_1 & A_1 \end{pmatrix} \begin{pmatrix} \bar{D}_1 & -\Lambda \bar{B}_1 \\ -\Lambda^{-1} \bar{C}_1 & \bar{A}_1 \end{pmatrix} \begin{pmatrix} A_1 & B_1 \\ C_1 & D_1 \end{pmatrix}, \\ \Phi_2 &:= \begin{pmatrix} \bar{D}_2 & -\bar{B}_2 \\ -\bar{C}_2 & \bar{A}_2 \end{pmatrix} \begin{pmatrix} A_2 & B_2 \\ C_2 & D_2 \end{pmatrix}. \end{aligned}$$

Proof. The loop γ_1 has four portions, as described above. The first portion is represented by the curve c_1 . The second portion is represented by $\kappa_3 \circ c_1^{-1}$. Using the facts that the third portion starts at the point $(1, \Lambda w_0)$ and that $\kappa_2 \circ \kappa_1 \circ \kappa_3(1, w_0) = (1, \Lambda w_0)$, we have that the third portion is represented by $\kappa_2 \circ \kappa_1 \circ \kappa_3 \circ c_1$. The final fourth portion is represented by $\kappa_1 \circ \kappa_2 \circ c_1^{-1}$, which follows from noting that $\kappa_1(1, w_0) = (1, w_0)$, $\kappa_2(1, w_0) = (1, w_0)$ and $\kappa_3(1, w_0) = (1, \Lambda w_0)$. (In particular, we see that $(1, w_0)$ is in the fixed point set of κ_1 and κ_2 but not in that of κ_3 .)

Thus we have that

$$\gamma_1 = (\kappa_1 \circ \kappa_2 \circ c_1^{-1}) \circ (\kappa_2 \circ \kappa_1 \circ \kappa_3 \circ c_1) \circ (\kappa_3 \circ c_1^{-1}) \circ c_1 .$$

Similarly, we can see that

$$\gamma_2 = (\kappa_1 \circ c_2^{-1}) \circ c_2 .$$

We can then apply Lemma 2.1 to get the result. $\square$

Using the Bryant type representation (1.3) to make CMC 1 surfaces in H^3 and S_1^3 , the conditions for the resulting surfaces to be well defined on M are that Φ_1 and Φ_2 are in $SU(2)$ and $SU(1, 1)$, respectively, and by symmetry, only the homotopy classes coming from γ_1 and γ_2 need be considered. However, the initial condition $F(1, w_0) = \text{id}$ will not cause Φ_1 and Φ_2 to lie in $SU(2)$ or $SU(1, 1)$. To remedy this, we will change the initial condition for the solution F so that it has initial condition

$$F(1, w_0) = \begin{pmatrix} \alpha & \beta \\ \beta & \alpha \end{pmatrix} \in \text{SL}(2, \mathbf{R})$$

in the case the ambient space is H^3 (that is, the $SU(2)$ case), and

$$F(1, w_0) = \begin{pmatrix} \alpha & \beta \\ \alpha & -\beta \end{pmatrix} \in \text{SL}(2, \mathbf{R})$$

in the case the ambient space is S_1^3 (that is, the $SU(1, 1)$ case).

Note that with these changes of initial condition of F , we still have enough symmetry to conclude that the resulting surfaces are well defined on M just by looking only at the two homotopy classes represented by γ_1 and γ_2 . This is because the homotopy group of M is generated by $[\gamma_1]$ and $[\gamma_2]$. The monodromies associated to those two homotopy classes are now, for $j = 1, 2$,

$$\begin{pmatrix} \alpha & \beta \\ \beta & \alpha \end{pmatrix}^{-1} \Phi_j \begin{pmatrix} \alpha & \beta \\ \beta & \alpha \end{pmatrix}$$

in the $SU(2)$ case, and

$$\begin{pmatrix} \alpha & \beta \\ \alpha & -\beta \end{pmatrix}^{-1} \Phi_j \begin{pmatrix} \alpha & \beta \\ \alpha & -\beta \end{pmatrix}$$

in the $SU(1, 1)$ case.

The closing conditions, that is, the conditions that the surfaces are well defined on M itself, are now that the above pairs of matrices lie in $SU(2)$ in the first case, and in $SU(1, 1)$ in the second case. Noting that Φ_1 and Φ_2 take the forms

$$\Phi_1 = \begin{pmatrix} r_1 & p \\ -\bar{p} & r_2 \end{pmatrix}, \quad \Phi_2 = \begin{pmatrix} q & ir_3 \\ ir_4 & \bar{q} \end{pmatrix}$$

for complex numbers p, q and real numbers r_1, r_2, r_3, r_4 , a direct computation gives that the closing conditions are

$$\frac{2 \operatorname{Re}(p)}{r_2 - r_1} = \frac{2 \operatorname{Im}(q)}{r_4 - r_3} = \frac{\alpha^2 + \beta^2}{2\alpha\beta} = \frac{1 + 2\beta^2}{2\beta\sqrt{1 + \beta^2}} \in (-\infty, -1) \cup (1, \infty)$$

in the $SU(2)$ case, and

$$\frac{2 \operatorname{Re}(p)}{r_2 - r_1} = \frac{2 \operatorname{Im}(q)}{r_4 - r_3} = \frac{\alpha^2 - \beta^2}{\alpha^2 + \beta^2} = \frac{1 - 4\beta^4}{1 + 4\beta^4} \in (-1, 1)$$

in the $SU(1, 1)$ case. So we have now proven the following lemma:

Lemma 2.3. *The single closing condition for one of the surfaces in (1.3) is that*

$$(2.2) \quad h_1(c, \lambda) = h_2(c, \lambda) \in \mathbf{R} \setminus \{\pm 1\},$$




FIGURE 4. A minimal surface in $\mathbf{R}^3$ (left) and a maximal surface in $\mathbf{R}_1^3$ (right), constructed as in Remark 2.4, with $k = 1$ and $\lambda = 2$, in each case showing closing with respect to one loop γ_j but not with respect to the other.

where

$$h_1(c, \lambda) = \frac{2 \operatorname{Re}(p)}{r_2 - r_1} \quad \text{and} \quad h_2(c, \lambda) = \frac{2 \operatorname{Im}(q)}{r_4 - r_3}$$

holds, and then the appropriate α and β can be found. Whether one obtains a surface in H^3 or S_1^3 is determined by whether the absolute value of $h_1(c, \lambda) = h_2(c, \lambda)$ is greater than or less than 1.

Remark 2.4. If we consider the minimal surface

$$\operatorname{Re} \int (1 - G^2, i(1 + G^2), 2G) \Omega \in \mathbf{R}^3$$

with the Weierstrass data (1.1), one can check that the period is solved for the loop γ_1 (regardless of the choice of λ , and we chose G as we did in (1.1) in order to make this true, see Remark 1.1), but is never solved for the loop γ_2 (for any choice of λ). On the other hand, if we consider the maximal surface

$$\operatorname{Re} \int (1 + G^2, i(1 - G^2), 2G) \Omega \in \mathbf{R}_1^3$$

with the Weierstrass data (1.1), the period is solved for γ_2 , but never for γ_1 . See Figure 4.

3. NUMERICAL EXPERIMENTS AND THE MAIN RESULT

Fix $\lambda = 2$. Here we provide constants $c \in \mathbf{R} \setminus \{0\}$ so that $h_1(c, 2) = h_2(c, 2)$ for the genus $k = 1, \dots, 20$ cases. It is a simple application of the intermediate value theorem to show $h_1(c, 2) = h_2(c, 2)$, and the functions $h_j(c, 2)$ are stable with respect to numerics if the paths $c_i(t)$ are chosen well – so the numerics are not delicate, and are expected to give reliable results. Furthermore, once we have existence of a surface for one value of $\lambda = \lambda_0$, we then have existence for all λ sufficiently close to λ_0 , so we can conclude existence of a 1-parameter family of such surfaces.

The data in Table 1 (see also Figure 5) then imply the numerical result stated in the introduction, except that we still need to analyze the behavior of the ends. Because $\deg(G) = k + 1$, equality in the Osserman inequality is satisfied, for both the H^3 and S_1^3 cases. From this, it follows that the ends are complete and embedded in the H^3 case, and also in the S_1^3 case when the ends are elliptic (see [RS] and

k	$c \in \mathbf{R} \setminus \{0\}$ so that $h_1(c, 2) =$ $h_2(c, 2)$	$h_j(c, 2)$	$c \in \mathbf{R} \setminus \{0\}$ so that $h_1(c, 2) =$ $h_2(c, 2)$	$h_j(c, 2)$	$c \in \mathbf{R} \setminus \{0\}$ so that $h_1(c, 2) =$ $h_2(c, 2)$	$h_j(c, 2)$
1	-0.0467552	-6.91432	-0.557726	0.130869	0.704094	0.221228
2	-0.0403901	-4.12613	-0.505010	0.218257	0.548964	0.0345248
3	-0.0334546	-3.32773	-0.483326	0.254392	0.482090	-0.0678105
4	-0.0281931	-2.95960	-0.471988	0.273656	0.444727	-0.132429
5	-0.0242574	-2.74968	-0.465097	0.285460	0.420845	-0.176931
6	-0.0212467	-2.61454	-0.460530	0.293371	0.404255	-0.209443
7	-0.0188836	-2.52044	-0.457291	0.299018	0.392055	-0.234233
8	-0.0169850	-2.45121	-0.454881	0.303237	0.382705	-0.253760
9	-0.0154287	-2.39818	-0.453020	0.306504	0.375309	-0.269538
10	-0.0141310	-2.35627	-0.451543	0.309105	0.369312	-0.282553
11	-0.0130330	-2.32232	-0.450342	0.311222	0.364352	-0.293472
12	-0.0120924	-2.29427	-0.449348	0.312979	0.360180	-0.302764
13	-0.0112778	-2.27070	-0.448511	0.314459	0.356623	-0.310766
14	-0.0105655	-2.25062	-0.447797	0.315722	0.353553	-0.317730
15	-0.00993749	-2.23331	-0.447182	0.316813	0.350878	-0.323846
16	-0.00937975	-2.21824	-0.446645	0.317764	0.348525	-0.329260
17	-0.00888113	-2.20499	-0.446173	0.318600	0.346439	-0.334086
18	-0.00843272	-2.19326	-0.445756	0.319341	0.344578	-0.338415
19	-0.00802733	-2.18280	-0.445383	0.320003	0.342907	-0.342319
20	-0.00765905	-2.17341	-0.445049	0.320596	0.341398	-0.345859

TABLE 1. Numerical results with $\lambda = 2$. The first column gives CMC 1 surfaces in H^3 . The second column gives CMC 1 faces in S_1^3 with elliptic ends. The third column gives CMC 1 faces in S_1^3 with hyperbolic ends.

[F2], for example). When the ends are hyperbolic in S_1^3 , they are neither complete nor embedded, but are weakly complete, because the metric (1.4) is complete, see [FRUY]. However, we have yet to show when the ends are elliptic or hyperbolic in the S_1^3 case. This final step is taken care of by the next lemma, which is similar to arguments found in the appendix of [F2], but here we are allowing for the case of general genus k .

Lemma 3.1. *The ends of the CMC 1 faces in S_1^3 in the middle column (resp. right hand column) of Table 1 have elliptic (resp. hyperbolic) ends.*

Proof. Let

$$F = \begin{pmatrix} A & B \\ C & D \end{pmatrix}$$

be a solution to (1.2). Note that, in order to determine the type of monodromy about an end, we only need to know the eigenvalues of the monodromy (provided those eigenvalues are not ± 1), and this is independent of the choice of F . So we may choose any solution to (1.2). We then have

$$X_{zz} + \left(\frac{1}{z} - \frac{w_z}{w} \right) X_z + \lambda^{k/(k+1)} \frac{cw_z}{zw} X = 0, \quad X = A, B,$$

and

$$Y_{zz} + \left(\frac{1}{z} + \frac{w_z}{w} \right) Y_z + \lambda^{k/(k+1)} \frac{cw_z}{zw} Y = 0, \quad Y = C, D.$$

Because of the symmetry $\kappa_2(z, w)$, it suffices to determine the type of just one end, and then the other end will automatically have the same type. So let us choose the




FIGURE 5. Graphs of the functions $h_1(c, 2)$ (thin curve) and $h_2(c, 2)$ (thick curve) in six cases: $k = 1$ in H^3 (upper left), $k = 1$ in S^3_1 with elliptic ends (upper middle), $k = 1$ in S^3_1 with hyperbolic ends (upper right), $k = 2$ in H^3 (lower left), $k = 2$ in S^3_1 with elliptic ends (lower middle), $k = 2$ in S^3_1 with hyperbolic ends (lower right).

end $(z, w) = (0, 0)$. At this end, w is a local coordinate for the Riemann surface $\overline{M}$. In terms of w , and considering z as a function of w , the equations above become

$$X_{ww} + o(1) \frac{X_w}{w} + \lambda^{k/(k+1)} c(k+1) (1 + o(1)) \frac{X}{w^2} = 0, \quad X = A, B,$$

and

$$Y_{ww} + 2(1 + o(1)) \frac{Y_w}{w} + \lambda^{k/(k+1)} c(k+1) (1 + o(1)) \frac{Y}{w^2} = 0, \quad Y = C, D,$$

where $o(1)$ denotes the Landau symbol, that is, $o(1)$ is a holomorphic function $\varphi(w)$ around $(z, w) = (0, 0)$ so that $\varphi(0) = 0$. It follows that the difference of the solutions of the indicial equation corresponding to the first of these two equations is

$$\sqrt{1 - 4c(k+1)\lambda^{k/(k+1)}}.$$

Likewise, the difference of solutions of the indicial equation for the second equation above takes the same value. It follows (see the appendix of [F2] for further details) that the end is elliptic (resp. hyperbolic) if

$$1 - 4c(k+1)\lambda^{k/(k+1)}$$

is positive (resp. negative) which is indeed the case for the data given in the middle column (resp. right hand column) of Table 1. $\square$

REFERENCES

- [AA] R. Aiyama and K. Akutagawa, *Kenmotsu-Bryant type representation formulas for constant mean curvature surfaces in $H^3(-c^2)$ and $S^3_1(c^2)$* , Ann. Global Anal. Geom. **17** (1998), 49-75.
- [B] R. Bryant, *Surfaces of mean curvature one in hyperbolic space*, Astérisque **154-155** (1987), 321-347.
- [F1] S. Fujimori, *Spacelike CMC 1 surfaces with elliptic ends in de Sitter 3-space*, Hokkaido J. Math. **35** (2006), 289-320.
- [F2] ———, *Spacelike mean curvature 1 surfaces of genus 1 with two ends in de Sitter 3-space*, Kyushu J. Math. **61** (2007), 1-20.
- [FRUY] S. Fujimori, W. Rossman, M. Umehara, K. Yamada and S-D. Yang, *Spacelike mean curvature one surfaces in de Sitter 3-space*, Comm. Anal. Geom. **17** (2009), 383-427.

- [FS] S. Fujimori and T. Shoda, *Minimal surfaces with two ends which have the least total absolute curvature*, preprint.
- [KY] Y. W. Kim and S-D. Yang, *A family of maximal surfaces in Lorentz-Minkowski three-space*, Proc. Amer. Math. Soc. **134** (2006), 3379–3390.
- [K] O. Kobayashi, *Maximal surfaces in the 3-dimensional Minkowski space L^3* , Tokyo J. Math. **6** (1983), 297–309.
- [LR] G. Levitt and H. Rosenberg, *Symmetry of constant mean curvature hypersurfaces in hyperbolic space*, Duke Math. J. **52** (1985), 53–59.
- [RS] W. Rossman and K. Sato, *Constant mean curvature surfaces in hyperbolic 3-space with two ends*, J. Exp. Math. **7** (1998), 101–119.
- [RUY1] W. Rossman, M. Umehara and K. Yamada, *Irreducible CMC- c surfaces in $H^3(-c^2)$ with positive genus*, Tohoku Math. J. **49** (1997), 449–484.
- [RUY2] ———, *Mean curvature 1 surfaces in hyperbolic 3-space with low total curvature*, Tohoku Math. J. **55** (2003), 375–395.
- [S] R. Schoen, *Uniqueness, symmetry, and embeddedness of minimal surfaces*, J. Diff. Geom. **18** (1983), 791–809.
- [UY1] M. Umehara and K. Yamada, *Complete Surfaces of Constant Mean Curvature One in the Hyperbolic 3-Space*, Ann. of Math., **137** (1993), 611–638.
- [UY2] ———, *Surfaces of Constant Mean Curvature c in $H^3(-c^2)$ with prescribed hyperbolic Gauss map*, Math. Ann. **304** (1996), 203–224.
- [UY3] ———, *A duality on CMC-1 surfaces in hyperbolic space, and a hyperbolic analogue of the Osserman inequality* Tsukuba J. Math. **21** (1997), 229–237.
- [UY4] ———, *Maximal surfaces with singularities in Minkowski space*, Hokkaido Math. J. **35** (2006), 13–40.

(Shoichi Fujimori) DEPARTMENT OF MATHEMATICS, FUKUOKA UNIVERSITY OF EDUCATION, MUNAKATA, FUKUOKA 811-4192, JAPAN

E-mail address: `fujimori@fukuoka-edu.ac.jp`

(Wayne Rossman) DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, KOBE UNIVERSITY, ROKKO, KOBE 657-8501, JAPAN

E-mail address: `wayne@math.kobe-u.ac.jp`